




PREPARATION FOR **MORNING PROCEDURE**

The bowel preparation induces frequent, loose bowel movements within one to three hours of taking the first dose. It is best to stay at home within easy reach of toilet facilities.

Please obtain from Chemist

2 x 15.5g PicoPrep sachets
1 x 70g Glycoprep-C sachet

2 Days Before		NO seeds or nuts
The Day Before	Breakfast & Lunch	<ul style="list-style-type: none"> Have light breakfast and light lunch. OK poached eggs, stewed fruit, clear soup, steamed chicken/fish NO seeds, no nuts, no bread, no cereals, no pasta, no rice After lunch have approved clear fluids for remainder of the day (refer to list of clear fluids below). NO solid food, no milk, no milk products <p>Mix the Glycoprep-C with 1 Litre of water and store in fridge for later. A small amount of clear (eg. lemon) cordial can be added to improve the taste.</p>
	At 3pm* 	<p>First Dose</p> <p>Mix 1 sachet PicoPrep with warm water (approx. 250ml). Stir until dissolved and allow to cool.</p> <ul style="list-style-type: none"> Drink mixture slowly and completely. Followed by a glass of approved clear fluids.
	At 6pm* 	<p>Second Dose</p> <ul style="list-style-type: none"> Drink the prepared Glycoprep-C slowly over 1 hour until finished. You may continue to drink approved clear fluids as required. <p>If feeling nausea or discomfort while drinking the preparation, simply slow down the rate of intake.</p>
	At 9pm 	<p>Third Dose</p> <p>Mix the last sachet of PicoPrep with warm water (approx. 250ml). Stir until dissolved and allow to cool.</p> <ul style="list-style-type: none"> Drink mixture slowly and completely; followed by a glass of approved clear fluid. Continue to drink at least 1 glass of approved clear fluid every hour until sleep.
Procedure Day		<ul style="list-style-type: none"> You should continue to drink approved clear fluids up to 6 hours prior to procedure. Nothing to eat or drink for the final 6 hours prior to procedure. You may take prescription medication (except diabetic medication) for up to 2 hours prior with a small sip of water.

* Alternate times for the day before procedure are: first dose 5pm, second dose 7pm and third dose 9pm.

Approved Clear Fluids	<p>water, clear fruit juice (apple, pear, grape), lemon jelly, black tea, black coffee, bonox, Lucozade, carbonated drinks, barley sugar, clear broth, lemon fruit cordial</p> <p>No red or purple colours</p>
------------------------------	---

Altona
Cnr Queen & Sargood St
Altona VIC 3018
T: 1800 4 HOBSON
F: 03 9398 8822

Sydenham
566 Melton Hwy
Sydenham VIC 3037
T: 1800 4 HOBSON
F: 03 8361 6700

Werribee
179 Princes Hwy
Werribee VIC 3030
T: 1800 4 HOBSON
F: 03 9731 8001